Troy Infusion Center 600 W Main Street Suite 120 Troy, OH 45373 Phone: 937-401-6620 Fax: 937-401-6629



Dayton, OH, 45459 Phone: 937-401-6620 Fax: 937-401-6629

Ferrlecit® (Ferric Gluconate) Order Form

Epic Referral Reference: REF133

| Patient Name: | DOB: | | | | |
|---|---|--|--|--|--|
| Address: | | | | | |
| Phone: | | | | | |
| ICD-10 Diagnosis Codes (2 required – 1 primary, 1 secondary): | | | | | |
| Primary Diagnosis Codes (pick one) | Secondary Diagnosis Codes (pick one) | | | | |
| □ D50.0 – Iron deficiency anemia secondary to blood loss | K90.9 – Intestinal malabsorption | | | | |
| □ D50.9 – Iron deficiency anemia, unspecified | □ K91.2 – Postsurgical malabsorption | | | | |
| □ D50.8 – Other iron deficiency anemias | □ T45.4X5D – Adverse effect of iron, subsequent encounter | | | | |
| □ O99.011 – Anemia complicating pregnancy 1 st trimester | □ Z87.19 – Personal history of other digestive disease | | | | |
| □ O99.012 – Anemia complicating pregnancy 2 nd trimester | | | | | |
| □ O99.013 – Anemia complicating pregnancy 3 rd trimester | | | | | |
| OR for Anemia related to chronic kidney disease: | | | | | |
| Primary Diagnosis Codes (pick one) | Secondary Diagnosis Codes (pick one) | | | | |
| □ N18.3 Chronic kidney disease, stage 3 (moderate) | \Box D50.0 – Iron deficiency anemia secondary to blood loss | | | | |
| □ N18.4 Chronic kidney disease, stage 4 (severe) | □ D50.8 – Other iron deficiency anemias | | | | |

□ D50.8 – Other iron deficiency anemias

□ D50.9 – Iron deficiency anemia, unspecified

□ D63.1 – Anemia in chronic kidney disease

Rx:

| Ferrlecit (ferric glucona | te) 125 mg in 100 mL 0 | .9% NaCl IV c | over 1 hour | | | |
|---------------------------|------------------------|---------------|-------------|-------|---------|--|
| Frequency: Daily | 2 times per week | Weekly | □ Every | weeks | Other _ | |

Total number of doses: _____

□ N18.5 Chronic kidney disease, stage 5

□ N18.6 End stage renal disease

Baseline labs must be included with the order (or available through Epic). Please note: follow-up iron labs should be completed \geq 4 weeks following last dose to evaluate full effect of iron repletion.

| **Port/PICC care per protocol will be performed | if applicable including heparin flush (500 units/5mL) and cathflo (2 mg) PRN | | | |
|---|--|--|--|--|
| for patients with a port** | | | | |
| Prescriber Printed Name: | | | | |
| Prescriber Full Address: | | | | |
| Office Phone Number: | Office Fax Number: | | | |
| Prescriber Signature: | Date: | | | |