

Troy Infusion Center  
600 W Main Street  
Suite 120  
Troy, OH 45373  
Phone: 937-401-6620  
Fax: 937-401-6629



Washington Township Infusion Center  
1989 Miamisburg-Centerville Road  
Suite 101  
Dayton, OH, 45459  
Phone: 937-401-6620  
Fax: 937-401-6629

## Injectafer® (Ferric Carboxymaltose) Order Form

Epic Referral: REF191

**Patient Name:** \_\_\_\_\_ **DOB:** \_\_\_\_\_

**Address:** \_\_\_\_\_

**Phone:** \_\_\_\_\_

**ICD-10 Diagnosis Codes (2 required – 1 primary, 1 secondary):**

<u>Primary Diagnosis Codes (pick one)</u>	<u>Secondary Diagnosis Codes (pick one)</u>
<input type="checkbox"/> D50.0 – Iron deficiency anemia secondary to blood loss	<input type="checkbox"/> K90.9 – Intestinal malabsorption
<input type="checkbox"/> D50.9 – Iron deficiency anemia, unspecified	<input type="checkbox"/> K91.2 – Postsurgical malabsorption
<input type="checkbox"/> D50.8 – Other iron deficiency anemias	<input type="checkbox"/> T45.4X5D – Adverse effect of iron, subsequent encounter
<input type="checkbox"/> O99.011 – Anemia complicating pregnancy 1 <sup>st</sup> trimester	<input type="checkbox"/> Z87.19 – Personal history of other digestive disease
<input type="checkbox"/> O99.012 – Anemia complicating pregnancy 2 <sup>nd</sup> trimester	
<input type="checkbox"/> O99.013 – Anemia complicating pregnancy 3 <sup>rd</sup> trimester	

**OR for Anemia related to chronic kidney disease:**

<u>Primary Diagnosis Codes (pick one)</u>	<u>Secondary Diagnosis Codes (pick one)</u>
<input type="checkbox"/> N18.3 Chronic kidney disease, stage 3 (moderate)	<input type="checkbox"/> D50.0 – Iron deficiency anemia secondary to blood loss
<input type="checkbox"/> N18.4 Chronic kidney disease, stage 4 (severe)	<input type="checkbox"/> D50.8 – Other iron deficiency anemias
<input type="checkbox"/> N18.5 Chronic kidney disease, stage 5	<input type="checkbox"/> D50.9 – Iron deficiency anemia, unspecified
<input type="checkbox"/> N18.6 End stage renal disease	<input type="checkbox"/> D63.1 – Anemia in chronic kidney disease

**Rx:**

Injectafer (ferric carboxymaltose) 750 mg in 250 mL normal saline infused IV over 20 minutes

- If patient weighs less than 50kg, give 15mg/kg instead of 750 mg.

**Frequency:**     Once         Weekly x 2 doses (doses must be at least 7 days apart)

*Baseline labs must be included with the order (or available through Epic). Please note: follow-up iron labs should be completed ≥ 4 weeks following last dose to evaluate full effect of iron repletion.*

\*\*Port/PICC care per protocol will be performed if applicable including heparin flush (500 units/5mL) and cathflo (2 mg) PRN for patients with a port\*\*

**Prescriber Printed Name:** \_\_\_\_\_

**Prescriber Full Address:** \_\_\_\_\_

**Office Phone Number:** \_\_\_\_\_ **Office Fax Number:** \_\_\_\_\_

**Prescriber Signature:** \_\_\_\_\_ **Date:** \_\_\_\_\_